

Operation of the *cbb*₃-Type Terminal Oxidase in *Azotobacter vinelandii*

Yu. V. Bertsova and A. V. Bogachev*

Department of Bioenergetics, Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Moscow, 119992 Russia; fax: (095) 939-0338; E-mail: sodiumc@genebee.msu.su

Received December 21, 2001

Revision received January 17, 2002

Abstract—A part of the gene encoding *cbb*₃-type cytochrome oxidase CcoN subunit was cloned from *Azotobacter vinelandii* and a mutant strain of this bacterium with disrupted *ccoN* gene was constructed. In contrast to the wild type strain, this one is unable to oxidize cytochromes *c*₄ and *c*₅. Thus, the *A. vinelandii* respiratory chain is shown to contain *cbb*₃-type cytochrome *c* oxidase. It is also shown that the activity of this enzyme is not necessary for diazotrophic growth of *A. vinelandii* at high oxygen concentrations.

Key words: *Azotobacter vinelandii*, cytochrome oxidase, *cbb*₃-type cytochrome, respiratory protection

A. vinelandii is a free-living, obligatory aerobic microorganism capable of nitrogen fixation. Most nitrogen-fixing bacteria are unable to reduce N₂ at high oxygen concentrations because the nitrogenase complex catalyzing this reaction is irreversibly inactivated by O₂. However, *A. vinelandii* fixes molecular nitrogen over a wide range of oxygen concentrations in spite of the fact that isolated nitrogenase complex of this bacterium is as sensitive to O₂ as that from other microorganisms [1]. To explain such oxygen-resistant nitrogen fixation of *Azotobacter*, Dalton and Postgate [2] advanced the hypothesis that the so-called respiratory protection mechanism functions in this bacterium. The term “respiratory protection” means a significant decrease in oxygen concentration in *Azotobacter* cytoplasm due to the active work of the respiratory chain to the level when nitrogenase inactivation does not occur.

The *A. vinelandii* respiratory chain includes at least two terminal oxidases, the *bd*-type quinol oxidase, and *o*-type cytochrome oxidase [3]. It was shown that mutations

in genes encoding cytochrome *bd* result in the loss of the unique ability of *A. vinelandii* for molecular nitrogen fixation at high O₂ concentrations [4]. Therefore, it was concluded that the respiratory protection of the nitrogenase complex is performed by the *bd*-type terminal oxidase activity [3, 4]. The operon encoding this enzyme was cloned and sequenced [4], and the enzyme itself was isolated and characterized [5, 6]. In particular, it was shown that activity of this terminal oxidase is coupled with proton potential generation, and the efficiency of this process is 1 H⁺/e⁻ [7].

In contrast to the *bd*-type quinol oxidase, the cytochrome oxidase branch of the *A. vinelandii* respiratory chain is so far poorly studied. It was demonstrated that the process of ubiquinol oxidation by this branch is sensitive to low concentrations of myxothiazol and antimycin [7]. Therefore, it was concluded that this process is realized via *bc*₁-complex. The presence of the following cytochromes *c*—*c*₄, *c*₅, *c*₅₅₁, and *c*₅₅₅—was demonstrated in *A. vinelandii* [8]. The last two types of cytochrome *c* are minor and their functions are unknown. As for cytochromes *c*₄ and *c*₅, their possible role is to maintain electron transfer from the *bc*₁-complex to cytochrome oxidase. A high TMPD-oxidase activity is typical of *A. vinelandii* mutant strains lacking genes encoding either cytochrome *c*₄ or cytochrome *c*₅, whereas the mutant strain lacking both these cytochromes *c* oxidizes TMPD very slowly [9, 10]. These data suggest that there are two alternative electron flows towards the cytochrome oxidase—via cytochrome *c*₄ as well as *c*₅ [9].

Abbreviations: PCR) polymerase chain reaction; SBP) sub-bacterial particles; TMPD) N,N,N',N'-tetramethyl-*p*-phenylenediamine; Ap^R) ampicillin resistance; Ap^S) sensitivity to ampicillin; H⁺/e⁻) number of protons transferred through the membrane by the respiratory chain enzyme normalized to the number of electrons; Km^R) kanamycin resistance; q/e⁻) charge number transferred through the membrane by the respiratory chain enzyme normalized to the number of electrons; Rf^R) rifampicin resistance; Tc^R) tetracycline resistance.

* To whom correspondence should be addressed.

Earlier the *o*-type terminal oxidase was isolated from *A. vinelandii* [11]. However, the N-terminal sequences in subunits were not determined in the course of this study, and the composition of cofactors of this enzyme also remained unknown. Thus, it remained unknown to what class of terminal oxidases does the *o*-type oxidase from *A. vinelandii* belongs. Later Leung and coauthors [12] cloned a part of the *A. vinelandii* gene (*cyoB*) probably encoding the *o*-type terminal oxidase. It was shown that this cloned fragment is homologous to the site of *cyoB* gene from *Escherichia coli* encoding the *bo*-type quinol oxidase subunit. Inactivation of this gene in *A. vinelandii* resulted in disappearance of the cell spectral characteristics ascribed to the *o*-type terminal oxidase. In this mutant strain CO binds only to cytochrome *bd* [12]. These data can also indicate that only two terminal oxidases are present in *A. vinelandii*. However, dependence of the respiration rate of *A. vinelandii* cells on oxygen concentration suggests that three terminal oxidases can function in this bacterium [3]. This suggestion is proved by the fact that along with genes encoding *bd*- and *o*-type terminal oxidases, *A. vinelandii* genome seems to contain also a gene homologous to *ccoN* gene of *cbb*₃-type cytochrome oxidase subunit [13]. The data can indicate that *cbb*₃-type enzyme exists in *A. vinelandii* as the third terminal oxidase, but this suggestion still needs experimental corroboration. Thus, the main goal of the present work was to study possible functioning of the *cbb*₃-type terminal oxidase in the *A. vinelandii* respiratory chain and to clarify the role of this enzyme in the respiratory protection mechanism.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *A. vinelandii* strains used in this work are listed in Table 1. *A. vinelandii* cells were grown on a thermostatted rotary shaker (250 rpm) at 30°C on modified Burck medium [7].

Sub-bacterial particles (SBP) were isolated from *A. vinelandii* cells as described earlier [14].

Cytochromes *c*₄ and *c*₅ were isolated from *A. vinelandii* as described in [15].

NADH oxidation by *A. vinelandii* SBP was measured using a Hitachi 557 spectrophotometer at 340 nm and 30°C. The measurement medium contained 20 mM Hepes (pH 7.5), 60 mM KCl, 2 mM MgSO₄, 0.9 μM gramicidin D, and SBP (1-5 μg protein/ml). To calculate the NADH oxidation rate, millimolar extinction coefficient $\epsilon_{340} = 6.22 \text{ mM}^{-1} \cdot \text{cm}^{-1}$ was used.

Oxidation of cytochromes *c* by *A. vinelandii* SBP was measured (at 30°C) using a Hitachi 557 spectrophotometer at 550-535 nm ($\epsilon_{550-535} = 17.3 \text{ mM}^{-1} \cdot \text{cm}^{-1}$) or at 420-440 nm ($\epsilon_{420-400} = 120 \text{ mM}^{-1} \cdot \text{cm}^{-1}$). The measurement medium contained 20 mM Hepes (pH 7.5), 60 mM KCl, 2 mM MgSO₄, reduced cytochrome *c* (final concentration of heme C was 5 μM) and SBP (10-50 μg protein/ml). Reduced cytochrome *c* was obtained by its incubation with ascorbate (5 mM) and subsequent purification from the reductant on a column with Sephadex G-25.

Spectral characteristics of SBP from various *A. vinelandii* strains were studied using an Aminco DW-2000 spectrophotometer. The measurement medium contained 100 mM KH₂PO₄, pH 7.0, and SBP (4-6 mg protein/ml). For complete oxidation of cytochromes, SBP were incubated in the presence of 1 mM potassium ferricyanide; to reduce cytochromes, a few crystals of sodium dithionite were added to the sample.

Protein concentration was measured by bicinchoninic acid method [16] using bovine serum albumin as the standard.

Construction of *A. vinelandii* strain with disrupted synthesis of CcoN protein. An *A. vinelandii* *ccoN* fragment was amplified by the polymerase chain reaction (PCR) method using RC3 5'-CGTCTGCGTCCGCTG and H6 5'-CCG-GAGTGGACGACGCCGATGGTCAG primers as described in [13]. An amplified fragment of *ccoN* gene was cloned into pGEM-T vector from Promega (USA) to obtain pC309 plasmid. Then a tetracycline resistance cassette from pHP45 ΩTc plasmid was incorporated into the unique *EcoRV* restriction site (positioned approximately 250 bp away from the 5'-end of PCR product). ΩTc-containing plasmid (pCT6) with the same direction of *ccoN* gene transcription and tetracycline resistance cassette was

Table 1. Bacterial strains and plasmids used in this work

Strain/plasmid	Genotype/phenotype	Reference
<i>A. vinelandii</i> UW136	Rf ^R	[12]
<i>A. vinelandii</i> DL10	UW136 <i>cyoB</i> ::Km, Rf ^R Km ^R	[12]
<i>A. vinelandii</i> C14	UW136 <i>ccoN</i> ::ΩTc, Rf ^R Tc ^R	this work
<i>A. vinelandii</i> DCO35	DL10 <i>ccoN</i> ::pCT6, Rf ^R Km ^R Tc ^R Ap ^R	same
pC309	pGEM-T with 900-bp <i>ccoN</i> fragment from <i>A. vinelandii</i> , Ap ^R	same
pCT6	pC309 with inclusion of ΩTc cassette in the <i>EcoRV</i> site of <i>ccoN</i> fragment, Ap ^R , Tc ^R	same

sampled. As a result of transformation of *A. vinelandii* UW136 cells by pCT6 plasmid, there was selected a C14 clone (*ccoN::ΩTc*) with Ap^STc^R phenotype typical of an insertion mutation obtained by a double homologous crossing-over event. Localization of mutation in *A. vinelandii* chromosome was proved by PCR method using RC3 primers (see above) and NC (5'-GAAG-GCCAGGGAGACCAC). Such PCR analysis of DNA from C14 clone revealed a unique 2.9-kb product (composed of 2.1- and 0.8-kb components corresponding to the tetracycline resistance cassette and PCR fragment of *ccoN* gene). We failed to detect the 0.8-kb PCR product typical of the wild type strain.

Mutagenesis of DL10 *A. vinelandii* cells (*cyoB::Km*) was performed analogously; a DCO35 strain (*cyoB::Km ccoN::pCT6*) in which *ccoN* inactivation occurred via the integration of pCT6 plasmid due to a single crossing-over, was thus obtained.

Competent *A. vinelandii* cells were obtained and transformed as described in [17].

Nucleotide sequencing was performed in the Human Genome Sequencing and Mapping Group of the Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, using an Applied Biosystems ABI 373A automatic sequencer.

The DNA sequence determined in this study was deposited into GenBank (No. AF456880).

RESULTS AND DISCUSSION

Study of the *cbb*₃-type cytochrome oxidase from *Rhodobacter capsulatus* demonstrated that gene fragment of this enzyme is also present in genomes of many other bacteria [13]. Among them, it was shown that *A. vinelandii* chromosome contains the *ccoN* fragment (encoding *cbb*₃-type cytochrome oxidase CcoN subunit in *R. capsulatus*) [13]. The data suggested that such terminal oxidase possibly functions also in *A. vinelandii*. To test this hypothesis, a mutant strain of *A. vinelandii* with disrupted synthesis of CcoN protein was constructed. For this, the *ccoN* gene fragment was amplified by PCR method as described in [13], and the products were separated by gel electrophoresis. The 900-bp band was isolated, cloned into pGEM-T vector (thus obtaining pC309 plasmid), and partially sequenced. Analysis of the resulting sequence using a BLASTX program showed that amino acid sequence corresponding to a certain nucleotide sequence is homologous to *cbb*₃-type terminal oxidase CcoN subunit from various bacteria. Maximal homology (99%) was found in CcoN protein analog from *Pseudomonas aeruginosa*. Thus, it was the *ccoN* gene fragment that we cloned from *A. vinelandii*. A partial nucleotide sequence of this gene was deposited into GenBank (No. AF456880).

The tetracycline resistance cassette was incorporated into the cloned fragment, and this resulted in pCT6 plasmid construction. To obtain the *A. vinelandii* strain with inactivated *ccoN* gene, cells of this bacterium were transformed by pCT6 plasmid (as all plasmids containing ColE1-replicon, this plasmid is unstable in *A. vinelandii* cells). A C14 clone with Ap^STc^R phenotype typical of mutation obtained due to a double crossing-over was sampled. Localization of the insertion mutation in *A. vinelandii ccoN* gene was proved by PCR analysis (see "Materials and Methods"). Thus, we constructed the *A. vinelandii* strain in which *ccoN* gene contains tetracycline resistance cassette; this should result in disruption of CcoN protein synthesis.

If *A. vinelandii* CcoN protein is a subunit of the functioning *cbb*₃-type terminal oxidase, then *ccoN* gene mutation should result in significant decrease in the cytochrome *c* oxidase activity in this microorganism, since all the so far described *cbb*₃-type oxidases are cytochrome oxidases. As shown in Fig. 1a, the dithionite-reduced *minus* air-oxidized difference spectrum of SBP isolated from the wild type *A. vinelandii* strain indicates the presence of cytochromes *c*, *b*, and *d* in membrane preparations of this bacterium. However, a peak at 553 nm is essentially absent from a similar (dithionite-reduced *minus* air-oxidized) difference spectrum of SBP from the CcoN-mutant *A. vinelandii* strain (C14); this may indicate a very low content of cytochromes *c* in these membrane preparations (Fig. 1a). Nonetheless, dithionite-reduced *minus* ferricyanide-oxidized difference spectra of SBP from these two strains are practically the same (Fig. 1b). The data mean that cytochromes *c* are present in SBP from the CcoN-mutant *A. vinelandii* strain, but in isolated SBP they are in reduced form, that is, in the mutant strain oxidation of cytochromes *c* by air oxygen is hindered, and they can be oxidized only in the presence of ferricyanide. The data indicate that the active cytochrome oxidase is absent from SBP from the *A. vinelandii* mutant strain.

To prove this suggestion, the fraction of cytochromes *c*₄ and *c*₅ was isolated from *A. vinelandii* cells, and the rate of oxidation of these cytochromes by SBP from various strains of the studied bacterium was measured. In accord with the previous observations [11], it was shown that SBP isolated from the *A. vinelandii* wild type strain (UW136) cells are able to oxidize cytochromes *c*₄ and *c*₅ (Table 2). The DL10 *A. vinelandii* strain mutant in the *o*-type terminal oxidase (*cyoB::Km*) demonstrated approximately the same cytochrome *c*₄/*c*₅ oxidase activity as the wild type strain (data not presented). The cytochrome *c* oxidase activity of these strains was completely inhibited in the presence of 20 μM KCN, whereas NADH oxidase activity was only partly sensitive (~15% inhibition) to such cyanide concentration, because NADH oxidase activity of *A. vinelandii* SBP is performed mainly due to the functioning of the *bd*-type "cyanide-resistant" quinol oxidase [7]. However, we failed to detect oxidation of




Fig. 1. Dithionite-reduced *minus* air-oxidized (a) and dithionite-reduced *minus* ferricyanide-oxidized (b) difference spectra of sub-bacterial particles isolated from the wild type *A. vinelandii* strain (1) and C14 mutant strain (2). All spectra are normalized to protein concentration 5 mg/ml.

cytochromes *c*₄ and *c*₅ by the CcoN-mutant *A. vinelandii* strain (C14), although the rate of NADH oxidation by SBP of this strain was approximately the same as that of the wild type strain (Table 2). The same results were obtained using cytochrome *c* from horse heart; the only difference was that the oxidation rate of this cytochrome was significantly lower compared with that of *A. vinelandii* cytochromes *c*₄ and *c*₅.

It is well known that NADH oxidation proceeds on the inner surface of the bacterial cytoplasmic membrane, whereas cytochrome *c* is oxidized on the outer surface of this membrane [18]. Thus, we had to ascertain that the ratio of “right-side out” and “inside out” particles in membrane preparations from the C14 mutant strain is not

changed compared with the wild type strain. As seen from the Table 2, stimulation of NADH oxidase activity by alamethicin was approximately the same in SBP from the mutant strain and in those from the wild type strain. Alamethicin is able to form channels of rather large size in the membrane in order to provide its permeability for NADH [19]. In the absence of alamethicin, only “inside out” particles catalyze NADH oxidation, whereas in the presence of this channel-former both populations of particles are catalysts (this stimulation of NADH oxidase activity by alamethicin cannot be rationalized by the effect of respiratory control, because all measurements were performed in the presence of gramicidin). Similar stimulation of NADH oxidase activity by alamethicin

Table 2. The oxidation rate of cytochromes *c*₄/*c*₅ and NADH by sub-bacterial particles from various *A. vinelandii* strains (results of a typical experiment are presented)

Strain	Activity			
	cytochrome <i>c</i> ₄ / <i>c</i> ₅ oxidase activity, nmol oxidized heme C/min per mg protein	NADH oxidase activity, nmol oxidized NADH/min per mg protein		
		– alamethicin	+ alamethicin*	ratio –/+ alamethicin**
UW136 (wild type)	33	11 900	16 000	74%
C14 (<i>ccoN</i> ::ΩTc)	0.0	8200	10 000	81%

* NADH oxidase activity in the presence of 17 μg/ml alamethicin.

** The ratio of NADH oxidase activity in the absence of alamethicin to this activity in the presence of alamethicin (%).




Fig. 2. Scheme of the terminal site of *A. vinelandii* respiratory chain.

(Table 2) means that orientation of SBP isolated from the *A. vinelandii* mutant strain does not differ from orientation of SBP from the wild type strain. Thus, the lack of cytochrome *c* oxidase activity in SBP from the C14 mutant strain can be explained only by the lack of the active cytochrome oxidase in this strain.

The data indicate that the *A. vinelandii* respiratory chain contains the active *cbb*₃-type cytochrome oxidase. Since *ccoN* gene mutation results in complete loss of cytochrome *c* oxidase activity, our data also indicate that all other terminal oxidases of *A. vinelandii* are quinol oxidases.

It should be noted that the oxidation rate of cytochromes *c* by SBP from *A. vinelandii* is significantly lower compared with the oxidation rate of NADH by the same SBP. In spite of the fact that the measured rate of cytochrome *c* oxidation is significantly underestimated due to a small content of "right-side out" particles in our membrane preparations (approximately fivefold, see Table 2), one can conclude that the cytochrome oxidase branch of the *A. vinelandii* respiratory chain is a minor component compared with the *bd*-type quinol oxidase.

As shown earlier, mutations in genes encoding the *bd*-type quinol oxidase result in disruption of diazotrophic growth of *A. vinelandii* at high oxygen concentrations [4]. However, the *A. vinelandii* strains with disrupted synthesis of *o*-type terminal oxidase are able to fix molecular nitrogen at high oxygen concentrations [12]. In this study we demonstrated that *ccoN* gene mutation (C14 strain) along with the double *ccoN cyoB* mutation (DCO35 strain) do not effect ability of *A. vinelandii* for diazotrophic growth at high oxygen concentrations (data not presented here). Thus, one can conclude that the *o*- and *cbb*₃-type terminal oxidases (in contrast to the *bd*-type quinol oxidase [4]) are not necessary for functioning of the respiratory protection mechanism of the nitrogenase complex.

Our results indicate that the *A. vinelandii* respiratory chain contains *cbb*₃-type cytochrome oxidase along with the *o*- and *bd*-type terminal oxidases [4, 12]. Accounting for the earlier results [4, 7, 10, 12], the terminal site of *A. vinelandii* respiratory chain can be presented as the following scheme (Fig. 2). The *bd*-type quinol oxidase possesses

low energy-conserving efficiency (1 q/e⁻) [7], a high rate of oxygen reduction and is a necessary component of the respiratory protection [4]. The *cbb*₃- and *o*-type terminal oxidases must possess high energy-conserving efficiency (2 q/e⁻, since they belong to superfamily of heme-copper oxidases [20]) and demonstrate relatively low activity with *A. vinelandii*. The physiological role of the two latter terminal oxidases seems to be energy supply of *A. vinelandii* cells during growth at low oxygen concentrations.

The authors are grateful to Professor R. Poole for a kind donation of *A. vinelandii* strains. This study was financially supported by the Russian Foundation for Basic Research (grant No. 01-04-48147) and special assistance for young scientists (grant No. 02-04-0610-mas).

REFERENCES

1. Yates, M. G. (1988) in *The Nitrogen and Sulphur Cycles* (Cole, I. A., and Ferguson, S. I., eds.) Cambridge, Cambridge University Press, Vol. 42, pp. 386-416.
2. Dalton, H., and Postgate, J. R. (1968) *J. Gen. Microbiol.*, **54**, 463-473.
3. Poole, R. K., and Hill, S. (1997) *Biosci. Rep.*, **17**, 303-317.
4. Kelly, M. J. S., Poole, R. K., Yates, M. G., and Kennedy, C. (1990) *J. Bacteriol.*, **172**, 6010-6019.
5. Junemann, S., and Wrigglesworth, J. M. (1995) *J. Biol. Chem.*, **270**, 16213-16220.
6. Kolonay, J. F., Moshiri, F., Gennis, R. B., Kaysser, T. M., and Maier, R. J. (1994) *J. Bacteriol.*, **176**, 4177-4181.
7. Bertsova, Y. V., Bogachev, A. V., and Skulachev, V. P. (1997) *FEBS Lett.*, **414**, 369-372.
8. Jones, C. W., and Redfearn, E. R. (1967) *Biochim. Biophys. Acta*, **143**, 340-353.
9. Ng, T. C. N., Laheri, A. N., and Maier, R. J. (1995) *Biochim. Biophys. Acta*, **1230**, 119-129.
10. Rey, L., and Maier, R. J. (1997) *J. Bacteriol.*, **179**, 7191-7196.
11. Jurtshuk, P., Mueller, T. J., and Wong, T. Y. (1981) *Biochim. Biophys. Acta*, **637**, 374-382.
12. Leung, D., van der Oost, J., Kelly, M., Saraste, M., Hill, S., and Poole, R. K. (1994) *FEMS Microbiol. Lett.*, **119**, 351-358.
13. Thöny-Meyer, L., Beck, C., Preisig, O., and Hennecke, H. (1994) *Mol. Microbiol.*, **14**, 705-716.
14. Bertsova, Y. V., Bogachev, A. V., and Skulachev, V. P. (1998) *Biochim. Biophys. Acta*, **1363**, 125-133.
15. Pettigrew, G. W., and Brown, K. R. (1988) *Biochem. J.*, **252**, 427-435.
16. Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., Fujimoto, E. K., Goeke, N. M., Olson, B. J., and Klenk, D. C. (1985) *Analyt. Biochem.*, **150**, 76-85.
17. Bertsova, Y. V., Bogachev, A. V., and Skulachev, V. P. (2001) *J. Bacteriol.*, **183**, 6869-6874.
18. Skulachev, V. P. (1988) *Membrane Bioenergetics*, Springer, Berlin.
19. Grivennikova, V. G., Kapustin, A. N., and Vinogradov, A. D. (2001) *J. Biol. Chem.*, **276**, 9038-9044.
20. Wikström, M., Bogachev, A., Finel, M., Morgan, J. E., Puustinen, A., Raitio, M., Verkhovskaya, M., and Verkhovskiy, M. I. (1994) *Biochim. Biophys. Acta*, **1187**, 106-111.