

Biochemical Characterization of Human Enteropeptidase Light Chain

M. E. Gasparian*, V. G. Ostapchenko, D. A. Dolgikh, and M. P. Kirpichnikov

*Shemyakin and Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, ul. Miklukho-Maklaya 16/10,
117997 Moscow, Russia; fax: (7-495) 336-8011; E-mail: marine_gasparian@yahoo.com*

Received July 7, 2005

Revision received October 17, 2005

Abstract—The synthetic gene encoding human enteropeptidase light chain (L-HEP) was cloned into plasmid pET-32a downstream from the gene of fusion partner thioredoxin immediately after the DNA sequence encoding the enteropeptidase recognition site. The fusion protein thioredoxin (Trx)/L-HEP was expressed in *Escherichia coli* BL21(DE3). Autocatalytic cleavage of the fusion protein and activation of recombinant L-HEP were achieved by solubilization of inclusion bodies and refolding of Trx/L-HEP fusion protein. The kinetic parameters of human and bovine enteropeptidases in the presence of different concentrations of Ca^{2+} and Na^{+} for cleavage of the specific substrate $\text{GD}_4\text{K-na}$ and nonspecific substrates such as small ester Z-Lys-SBzl and chromogenic substrates Z-Ala-X-Arg-pNA have been comparatively analyzed. It is demonstrated that positively charged ions increased the Michaelis constant (K_m) for cleavage of specific substrate $\text{GD}_4\text{K-na}$, while the catalytic constant (k_{cat}) remained practically unchanged. L-HEP demonstrated secondary specificity to the chromogenic substrate Z-Ala-Phe-Arg-pNA with k_{cat}/K_m 260 $\text{mM}^{-1}\cdot\text{sec}^{-1}$. Enzymatic activity of L-HEP was suppressed by inhibitors of trypsin-like and cysteine (E-64), but not metallo-, amino-, or chymotrypsin-like proteinases. L-HEP was active over a broad range of pH (6-9) with optimum activity at pH 7.5, and it demonstrated high stability to different denaturing agents.

DOI: 10.1134/S0006297906020015

Key words: human enteropeptidase, recombinant fusion proteins, specificity, kinetic parameters, protease inhibitors

Enteropeptidase (synonym: enterokinase) (EC 3.4.21.9) is a serine protease of the intestinal brush border that activates trypsinogen by cleavage of the N-terminal peptide following the sequence $(\text{Asp})_4\text{-Lys}$. As a result, active trypsin is generated which activates many pancreatic zymogens [1]. Native dimer of enteropeptidase has been purified from porcine [2], bovine [3, 4], and human [5] intestine. In all cases, enteropeptidase appears to be a

disulfide-linked heterodimer that is derived from a single-chain precursor; it is composed of the heavy (82-140 kD) and the light (35-62 kD) chains. The light chain is a catalytic subunit of enteropeptidase that consists of a chymotrypsin-like serine protease domain. Human enteropeptidase mRNA expression is limited to the proximal small intestine, and the protein is found in enterocytes of the duodenum and proximal jejunum [6, 7]. Congenital deficiency of enteropeptidase causes severe intestinal malabsorption with diarrhea, vomiting, and growth failure. These abnormalities can be treated successfully by supplementation with pancreatic extract [8, 9]. Mutations in the proenteropeptidase gene are the molecular cause of congenital enteropeptidase deficiency [10]. The high degree of specificity exhibited by enteropeptidase makes it an ideal reagent for cleaving bacterially produced fusion proteins. The expression and purification of the catalytic subunit of bovine enteropeptidase from *E. coli* [11], the methylotrophic yeast *Pichia pastoris* [12], the filamentous fungus *Aspergillus niger* [13], and *Saccharomyces cerevisiae* [14] have been described.

Abbreviations: L-HEP) recombinant human enteropeptidase light chain; L-BEP) recombinant bovine enteropeptidase light chain purified from *E. coli*; Trx/L-HEP) thioredoxin/human enteropeptidase light chain fusion protein; Trx/hEGF) thioredoxin/human epidermal growth factor fusion protein; Z-Lys-SBzl) thiobenzyl benzyloxy-carbonyl-L-lysinate; $\text{GD}_4\text{K-na}$) Gly-Asp-Asp-Asp-Asp-Lys- β -naphthylamide; Z-Ala-X-Arg-pNA) benzyloxycarbonyl-L-alanine-X-L-arginine-*p*-nitroanilide; STI) soybean trypsin inhibitor; DTNB) 5,5'-dithiobis-(2-nitrobenzoic acid); TPCK) N-tosyl-L-phenylalanine chloromethyl ketone; TLCK) N-tosyl-L-lysine chloromethyl ketone.

* To whom correspondence should be addressed.

In all cases, the recombinant proteins demonstrated high and remarkably specific protease activity. Kinetic parameters of wild type and various mutant forms have been determined, and this revealed a critical substrate-binding segment outside the active site [15]. Finally, the crystal structure of bovine enteropeptidase light chain complexed with an analog of the trypsinogen-activation peptide has been solved to 2.3 Å resolution [15].

In this paper, we report cloning and expression in *E. coli* of the synthetic gene encoding the human enteropeptidase catalytic subunit with subsequent refolding and purification. The influence of ionic strength, pH, different denaturing agents, and protease inhibitors has been investigated for the cleavage of synthetic substrates as well as of fusion proteins by L-HEP (recombinant human enteropeptidase light chain) and L-BEP (recombinant bovine enteropeptidase light chain purified from *E. coli*).

MATERIALS AND METHODS

Materials. Host strain *E. coli* BL21(DE3) and plasmid pET-32a were from Novagen (USA). GD₄K-na (Gly-Asp-Asp-Asp-Asp-Lys-β-naphthylamide), Z-Lys-SBzl (thiobenzyl benzyloxy-carbonyl-L-lysinate), STI (soybean trypsin inhibitor), aprotinin, leupeptin, antipain, E-64, benzamidin hydrochloride, chymostatin, pepstatin A, bestatin, TPCK (N-tosyl-L-phenylalanine chloromethyl ketone), TLCK (N-tosyl-L-lysine chloromethyl ketone), STI-agarose, and DTNB (5,5'-dithiobis-(2-nitrobenzoic acid)) were from Sigma (USA). Recombinant bovine enteropeptidase light chain, restriction enzymes, and DNA polymerase were from New England Biolabs (USA).

Synthesis of the L-HEP gene and its cloning into pET-32a vector. The DNA sequence encoding the light chain of human enteropeptidase (705 bp) was synthesized from 26 oligonucleotides (35-43 nucleotides each) by polymerase chain reaction (PCR) as described before [16]. The amplified DNA sequence was cleaved by *Bgl*II and *Hind*III and cloned into plasmid pET-32a at *Bgl*II/*Hind*III restriction sites downstream to the gene of the fusion partner thioredoxin following the DNA sequence encoding (Asp)₄-Lys. The L-HEP gene sequence was confirmed by automatic sequencing on an ABI Prism-310 Genetic Analyzer (Applied Biosystems, USA).

Expression, renaturation, and purification of L-HEP. *Escherichia coli* BL21(DE3) cells were transformed by plasmid pET-32a/L-HEP. Expression, renaturation, and purification of L-HEP were done as described in our previous paper [16]. Total autocatalytic cleavage of the Trx/L-HEP (thioredoxin/human enteropeptidase light chain fusion protein) was observed after 2 h dialysis against 0.1 M Tris-HCl (pH 8.0) containing 1 mM CaCl₂ at room temperature. NaCl to the final concentration of

500 mM was added to the dialyzed protein, and the active enzyme was purified by affinity chromatography on STI-agarose. The purified protein was stored in buffer containing 0.1 M Tris-HCl (pH 8.0), 50% glycerol, and 500 mM NaCl at -20°C. The concentration of L-HEP was determined spectrophotometrically by absorbance at 280 nm using a molar extinction coefficient of 54,450 M⁻¹·cm⁻¹ calculated from the known amino acid sequence using the ProtParam tool on the ExPASy server (www.expasy.org).

The rate of hydrolysis of the fluorogenic substrate GD₄K-na was determined as described previously [17]. Assays were performed at 37°C in 1.5 ml buffer containing 0.01-1.0 mM GD₄K-na, 25 mM Tris-HCl (pH 8.4), and 10% dimethylsulfoxide with different concentrations of CaCl₂. The reaction was initiated by adding enteropeptidase (final concentration 0.7 nM). The rate of β-naphthylamine formation was determined from the increment of fluorescence ($\lambda_{\text{ex}} = 337$ nm and $\lambda_{\text{em}} = 420$ nm) monitored continuously for 5 min using MPF-4 fluorimeter (Hitachi, Japan).

Rates of hydrolysis of synthetic chromogenic tripeptides Z-Ala-X-Arg-pNA (benzyloxycarbonyl-L-alanine-X-L-arginine-*p*-nitroanilide) were measured in 1 ml buffer containing 0.1 M Tris-HCl (pH 8.0), 5% dimethylsulfoxide, with concentration range of the substrates 0.01-1 mM. Reactions were started by addition of L-HEP to the final concentration of 4 nM, and absorbance of the released *p*-nitroaniline was recorded at 405 nm continuously for 5 min. The k_{cat} and K_{m} values for all substrates were determined from Lineweaver-Burk plots.

Effects of protease inhibitors and denaturing reagents on L-HEP activity. The reaction of hydrolysis of thiobenzyl benzyloxycarbonyl-L-lysinate (Z-Lys-SBzl) was used for measurements of L-HEP activity in the presence of protease inhibitors and denaturing reagents as described previously [18]. Assays were performed at room temperature in 1.5 ml buffer containing 0.1 M Tris-HCl (pH 8.0), 260 μM DTNB, and 66 μM Z-Lys-SBzl. After addition of the enzyme (final concentration 0.5 nM) the rate of 3-carboxy-4-nitrothiophenoxide production was measured continuously for 5-30 min from the absorbance changes at 412 nm, using the extinction coefficient 13,600 M⁻¹·cm⁻¹.

Cleavage of fusion proteins by L-HEP and L-BEP. Trx/hEGF (thioredoxin/human epidermal growth factor fusion protein) was partially purified from inclusion bodies by affinity chromatography on Ni-NTA agarose. Digestions by L-HEP and L-BEP were held in buffer containing 50 mM Tris-HCl (pH 8.0) in the presence of different denaturing agents during 20 h at room temperature. After digestion, samples were analyzed on Tris-Tricine SDS-PAGE [19]. Protein electrophoretic profiles in gels were stained with Coomassie Blue G-250, and the bands were scanned using an ImageMaster VDS densitometer (Amersham Pharmacia Biotech, USA). The

Trx/hEGF concentration was determined by the Bradford method [20].

RESULTS AND DISCUSSION

L-HEP gene expression in *E. coli* and L-HEP purification. The DNA sequence encoding the light chain of human enteropeptidase (GenBank Accession Number U09860) was synthesized from 26 oligonucleotides by PCR and cloned into plasmid pET-32a after the DNA sequence encoding enteropeptidase recognition site [16]. High level expression of Trx/L-HEP (100 mg from 100 ml culture) was achieved in *E. coli* strain BL21(DE3). Most (90%) of the fusion protein was insoluble and formed inclusion bodies. After solubilization of inclusion bodies in 6 M guanidine-HCl, Trx/L-HEP fusion protein was refolded using oxidized and reduced glutathione as described in our previous work [16]. Refolded protein cleaved autocatalytically. Active enzyme was purified by affinity chromatography on STI-agarose. This approach

yielded more than 1 mg of highly active enzyme from 100 ml of *E. coli* culture (Table 1).

Estimation of L-HEP and L-BEP kinetic parameters for cleavage of the specific substrate GD₄K-na at different concentrations of Ca²⁺ and Na⁺. Conventionally, kinetic parameters for the specific substrate GD₄K-na cleavage by bovine enteropeptidase light chain have been determined in the presence of 10 mM CaCl₂ [15, 21]. Our measurements have shown that the maximum cleavage rate is observed at 0.02 mM CaCl₂ both for L-HEP and L-BEP with the specificity constants (k_{cat}/K_m) of 3464 and 347 mM⁻¹·sec⁻¹, respectively. For hydrolysis of the substrate at 10 mM CaCl₂, 4.5-fold lower values of specificity constants were observed (Table 2, Fig. 1a). It has been shown that Ca²⁺ mainly increases K_m , while k_{cat} remains practically unchanged. The rate of GD₄K-na hydrolysis was inhibited with 100 and 250 mM NaCl by 70 and 90%, respectively (data not shown). Similar results were obtained for cleavage of Trx/hEGF fusion at different concentrations of CaCl₂ (Fig. 1b). Even 1 mM CaCl₂ slightly diminished the rate of Trx/hEGF hydrolysis, while 5 and 10 mM CaCl₂ reduced it by 50 and 80%, respectively. The absence of Ca²⁺ stimulation of enteropeptidase activity is consistent with Leu and Thr at positions 70 and 80 in the polypeptide chain (chymotrypsin numbering), while Glu at these positions are involved in formation of a Ca²⁺ binding site in thrombin and trypsin [22].

However, addition of 10 mM CaCl₂ had no effect on the rate of hydrolysis of nonspecific substrates Z-Lys-SBzl and Z-Ala-X-Phe-Arg-pNA (data not shown), and NaCl at very high concentration (up to 1-5 M) increased (2-2.5-fold) the rate of their hydrolysis by the enteropeptidase (Table 3). This suggests that Ca²⁺ or Na⁺ successfully shields substrate (Asp)₄ negatively charged side chains thereby impairing its binding to the positively charged substrate-binding RRRK site of the enzyme. One

Table 1. Yield of purified active L-HEP after refolding of Trx/L-HEP fusion protein

Fusion protein	Inclusion bodies before renaturation from 100 ml cell culture, mg	Refolded and purified on STI-agarose active enteropeptidase, mg	Yield of refolding, %
Trx/L-HEP	100 (58.6 mg L-HEP)	1.2 ± 0.2	2.0

Note: Values are from five independent experiments.

Table 2. Kinetic parameters for the cleavage of the specific substrate GD₄K-na by L-HEP and L-BEP

Enteropeptidase	[EDTA], mM	[CaCl ₂], mM	K_m , mM	k_{cat} , sec ⁻¹	k_{cat}/K_m , mM ⁻¹ ·sec ⁻¹
L-HEP	1.0		0.080 ± 0.004	117 ± 5	1462
L-BEP	1.0		0.332 ± 0.031	46.0 ± 2.1	138
L-HEP		0.02	0.034 ± 0.002	118 ± 5	3464
L-BEP		0.02	0.141 ± 0.023	49.3 ± 2.1	347
Human dimer*		0.1	0.525	28.3	41
L-HEP		0.1	0.042 ± 0.003	117 ± 5	2600
L-BEP		0.1	0.189 ± 0.007	46.6 ± 2.2	243
Human dimer*		10	0.28	21.5	101
L-HEP		10	0.160 ± 0.007	115 ± 4	719
L-BEP		10	0.590 ± 0.031	45.4 ± 2.3	78

Note: Values for K_m and k_{cat} are expressed as mean ± S.D. of three independent determinations.

* Data from [15].




Fig. 1. Effect of CaCl_2 on the hydrolysis rate of the GD_4K -na peptide and Trx/hEGF fusion protein cleavage by L-HEP and L-BEP. a) The rate of GD_4K -na hydrolysis at different concentrations of CaCl_2 or 1 mM EDTA (point 0 mM in the diagram). b, c) Separate samples containing 15 μg of Trx/hEGF fusion were incubated with the same relative activity units of enzymes (L-HEP and L-BEP) for 16 h at 25°C in 0.1 M Tris-HCl, pH 8.0, at different concentrations of CaCl_2 (b) or at 0.02 mM CaCl_2 for various time intervals (c). Samples were analyzed by 12% Tris-Tricine SDS-PAGE and the gel was stained with Coomassie brilliant blue.

Table 3. Influence of different denaturing agents on the rate of Z-Lys-SBzl hydrolysis by L-HEP

Denaturant	Concentration of denaturant, %	Activity, %	Denaturant	Concentration of denaturant, M	Activity, %
Triton X-100	0.1	70 ± 3	Imidazole	1.5 · 10 ⁻¹	87 ± 3
	0.5	30 ± 1		2.5 · 10 ⁻¹	68 ± 2
	1.0	18 ± 1	DTT	1.0 · 10 ⁻²	100 ± 3
Tween-20	0.1	80 ± 4		2.0 · 10 ⁻²	91 ± 3
	0.5	35 ± 2	NaCl	1.0 · 10 ⁻¹	150 ± 7
	1.0	25 ± 1		1.0	225 ± 10
SDS	0.1	60 ± 3		3.0	250 ± 10
	0.5	35 ± 2	5.0	145 ± 7	
	1.0	28 ± 1	KCl	1.0 · 10 ⁻¹	120 ± 6
Ethanol	5.0	79 ± 4		1.0	100 ± 5
	10	70 ± 3		3.0	100 ± 5
	Urea	20	62 ± 3	1.0	50 ± 3
				2.0	30 ± 2

Note: Data are presented as means ± S.D. from at least three independent experiments.

Table 4. Kinetic parameters for the cleavage of nonspecific substrates by L-HEP

Substrate	k_{cat} , sec ⁻¹	K_m , μ M	k_{cat}/K_m , mM ⁻¹ .sec ⁻¹
Z-Lys-SBzl	133 ± 4	140 ± 7	950 ± 30
For-Ala-Phe-Arg-pNA	23.5 ± 0.9	76 ± 4	309 ± 11
Z-Ala-Phe-Arg-pNA	17.5 ± 0.7	67 ± 4	261 ± 10
Z-Gly-Pro-Arg-pNA	14.4 ± 0.6	91 ± 5	158 ± 7
For-Ala-Phe-Lys-pNA	18.2 ± 0.8	167 ± 8	109 ± 5
Z-Ala-Tyr-Arg-pNA	10.0 ± 0.4	109 ± 5	92 ± 5
Z-Ala-Leu-Arg-pNA	15.2 ± 0.7	234 ± 11	65 ± 3
Z-Ala-Trp-Arg-pNA	11.5 ± 0.5	240 ± 12	48 ± 3
Z-Ala-Ala-Arg-pNA	9.5 ± 0.7	356 ± 16	26.7 ± 1.6
Z-Pro-Phe-Arg-pNA	6.2 ± 0.3	265 ± 13	23.4 ± 1.3
Z-Ala-Met-Arg-pNA	7.6 ± 0.3	420 ± 20	18.2 ± 1.0

Note: Data are presented as mean ± S.D. from at least three independent experiments.

should note that despite high amino acid sequence homology of human and bovine enteropeptidase light chains (84%), L-HEP cleaved the specific substrate GD₄K-na almost 10 times faster than L-BEP at any concentration of CaCl₂.

In our previous work, we have shown that the kinetic parameters for cleavage of the small ester Z-Lys-SBzl by human and bovine enzymes are practically identical [16]. This observation let us carry out a comparative analysis of cleavage rates of different fusion proteins, in particular, Trx/hEGF by L-HEP and L-BEP (Fig. 1c). After quantification of Trx/hEGF bands on SDS-PAGE gels by densitometry, the rates of hydrolysis of Trx/hEGF fusion by L-HEP and L-BEP (2.1 and 0.22 mmol/min per mg, respectively) were calculated. Additional studies with site-directed mutagenesis are required to explain higher (about an order of magnitude) activity of L-HEP in comparison with L-BEP.

Secondary substrate specificity of L-HEP. Kinetic parameters of L-HEP were determined for hydrolysis of the thioester Z-Lys-SBzl and some synthetic tripeptides (Table 4). Kinetic parameters for thioester Z-Lys-SBzl cleavage by L-HEP and L-BEP were practically identical. Hydrolysis of the tripeptide substrates For-Ala-Phe-Arg-pNA and For-Ala-Phe-Lys-pNA indicated a 2.5-fold Arg to Lys preference in P1 position. L-HEP demonstrated relatively high affinity to chromogenic tripeptides with hydrophobic residues in the P2 position (Phe > Tyr > Leu > Trp). For example, Z-Ala-Phe-Arg-pNA was hydrolyzed with quite high specificity constants k_{cat}/K_m 260 and 140 mM⁻¹.sec⁻¹ by L-HEP and L-BEP, respectively. We believe that a substrate with even greater specificity constant can be obtained by optimizing amino acids at positions P3 and P4 of the substrate. Similar value of

$k_{cat}/K_m = 212 \text{ mM}^{-1} \cdot \text{sec}^{-1}$ was obtained for native porcine enteropeptidase toward Z-Phe-Arg-MCA [23].

Calcium ions (up to 20 mM) had no effect on hydrolysis rate of Z-Lys-SBzl and tripeptides, and NaCl (1–3 M) only stimulated 2.0–2.5 times the rate of hydrolysis (Table 3). All hereinabove data indicated that the binding site for Z-Ala-X-Arg-pNA tripeptides differs from that for the specific substrate GD₄K-na. It remains possible that other targets playing an important physiological role may exist for the human enteropeptidase proteolytic domain under certain conditions.

Interestingly, in the phylogenetic tree of 84 serine proteases of the chymotrypsin family enteropeptidase was found in one group with hepsin, plasma kallikrein, prostaticin, and factor XI, which prefer Arg and Phe at the P1 and P2 positions of substrates, respectively [22]. This suggests that P1Arg-P2Phe secondary substrate specificity was the original one for enteropeptidase during evolution. Detailed analysis of the evolution of trypsinogen activation peptide demonstrated that the (Asp)₄-Lys sequence in mammalian trypsinogen has evolved to inhibit autoactivation and enhance cleavage by enteropeptidase [24].

Influence of denaturing agents on the enzymatic activity of L-HEP. The stability of L-HEP to denaturing




Fig. 2. Effect of different denaturing agents on the cleavage rate of Trx/hEGF fusion protein by L-HEP. Samples containing 15 μ g of Trx/hEGF fusion were incubated for 20 h at 25°C in buffer containing 0.1 M Tris-HCl (pH 8.0) and different concentrations of imidazole, NaCl, or Triton X-100.

agents was examined using Z-Lys-SBzl (Table 3) and Trx/hEGF fusion protein (Fig. 2) as substrates. The enzyme appeared highly resistant to detergents like Triton X-100, Tween 20, and SDS. L-HEP hydrolyzed Z-Lys-SBzl in 20% ethanol preserving about 60% of activity. High concentrations of imidazole (250 mM) and dithiothreitol (20 mM), usually used in purification of fusion proteins, were acceptable for cleavage of both Z-Lys-SBzl and Trx/hEGF fusion proteins. High concentrations of NaCl (up to 5 M) or KCl (up to 3 M) only stimulated cleavage of the nonspecific substrates Z-Lys-SBzl and Z-Ala-Phe-Arg-pNA, while cleavage of the specific substrate GD₄K-na was strongly inhibited even by low concentrations of NaCl (100 mM and more, data not shown).

Effect of protease inhibitors on L-HEP. The effect of various protease inhibitors on L-HEP was studied using the thioester Z-Lys-SBzl as a substrate (Table 5). The activity of enteropeptidase was strongly inhibited by trypsin-like serine protease inhibitors. STI and aprotinin inhibited the enzymes in a time-dependent manner (with equilibrium established in 30–60 min), while leupeptin,

Table 5. Effects of protease inhibitors on hydrolytic activity of L-HEP

Inhibitor	Concentration of inhibitor, M	Activity of L-HEP, %
STI	$3.0 \cdot 10^{-9}$	46.1 ± 1.1
	$3.3 \cdot 10^{-8}$	6.0 ± 0.2
Aprotinin	$2.5 \cdot 10^{-7}$	36.0 ± 1.0
	$2.0 \cdot 10^{-6}$	8.8 ± 0.3
Leupeptin	$1.0 \cdot 10^{-5}$	6.3 ± 0.3
Antipain	$1.0 \cdot 10^{-5}$	15.7 ± 1.0
	$1.0 \cdot 10^{-4}$	4.5 ± 0.2
E-64	$1.0 \cdot 10^{-5}$	52.4 ± 1.4
	$1.0 \cdot 10^{-4}$	16.3 ± 0.8
Benzamidine	$1.0 \cdot 10^{-4}$	35.6 ± 0.9
	$1.0 \cdot 10^{-3}$	8.2 ± 0.5
Chymostatin	$1.0 \cdot 10^{-4}$	81.1 ± 2.0
Pepstatin A	$1.0 \cdot 10^{-4}$	95.0 ± 1.5
Bestatin	$1.0 \cdot 10^{-4}$	93.0 ± 1.8
PMSF	$1.0 \cdot 10^{-3}$	45.9 ± 1.1
TLCK	$1.0 \cdot 10^{-3}$	87.0 ± 1.5
EDTA	$1.0 \cdot 10^{-2}$	100.0 ± 3.3

Note: Data are presented as mean \pm S.D. from at least three independent experiments.




Fig. 3. The pH dependence of Z-Lys-SBzl hydrolysis by L-HEP. Data (filled squares) were treated to fit a Gaussian curve with pH optimum 7.54 ± 0.02 .

antipain, and benzamidine did it instantaneously. L-HEP activity was inhibited very poorly by TLCK and was practically not influenced by chymotrypsin-like protease (chymostatin), aminoprotease (bestatin), metalloprotease (EDTA), or aspartic protease (pepstatin) inhibitors. Surprisingly, cysteine protease inhibitor E-64 inhibited L-HEP efficiently. Earlier the same inhibition by E-64 was demonstrated for porcine enteropeptidase [23] and β -trypsin [25]. It has been shown by computer modeling that inhibition of β -trypsin was particularly determined by the interaction of the guanidine-terminal group of E-64 with S1-Asp189 side chain. In contrast, in cysteine proteinases the epoxide ring from the opposite terminus of E-64 reacts with the catalytic site thiolate anion [26]. Inhibition of enteropeptidase can be additionally related to electrostatic interaction of the carboxy-terminal group of E-64 with the 96–99 (RRRK) loop of L-HEP.

The pH dependence of L-HEP enzymatic activity was measured for hydrolysis of Z-Lys-SBzl as described in “Materials and Methods”. The enzyme cleaved it over a broad pH range with pH optimum 7.5 and 50% activity at pH 6.3 and 8.7 (Fig. 3).

Thus, we have obtained and characterized for the first time the human enteropeptidase catalytic subunit. Our preparation retained 100% activity during storage at -20°C for at least for 1.5 years. High activity, specificity, and stability of human enteropeptidase make it a very useful reagent in biotechnology for the cleavage of fusion proteins.

We thank Anna Gorozhanina, a student from the Moscow State University Department of Bioengineering, for skilful technical assistance.

The study was supported by the Russian Academy of Sciences Program on Molecular and Cell Biology.

REFERENCES

1. Kunitz, M. (1939) *J. Gen. Physiol.*, **22**, 429-446.
2. Baratti, J., Maroux, S., Louvard, D., and Desnuelle, P. (1973) *Biochim. Biophys. Acta*, **315**, 147-161.
3. Anderson, L. E., Walsh, K. A., and Neurath, H. (1977) *Biochemistry*, **16**, 3354-3360.
4. Liepnieks, J. J., and Light, A. (1979) *J. Biol. Chem.*, **254**, 1677-1683.
5. Magee, A. I., Grant, D. A., and Taylor, J. H. (1981) *Clin. Chim. Acta*, **115**, 241-254.
6. Lojda, Z., and Gossrau, R. (1983) *Histochemistry*, **78**, 251-270.
7. Mitoshi, Y., Onishi, T., Sano, T., and Komi, N. (1990) *Gastroenterol. Jpn.*, **25**, 320-327.
8. Hadorn, B., Tarlow, M. J., Lloyd, J. K., and Wolff, O. H. (1969) *Lancet*, **1**, 812-813.
9. Haworth, J. C., Gourley, B., Hadorn, B., and Sumida, C. (1971) *J. Pediatr.*, **78**, 481-490.
10. Holzinger, A., Maier, E. M., Buck, C., Mayerhofer, P. U., Kappler, M., Haworth, J. C., Moroz, S. P., Hadorn, H. B., Sadler, J. E., and Roscher, A. A. (2002) *Am. J. Hum. Genet.*, **70**, 20-25.
11. Collins-Racie, L. A., McColgan, J. M., Grant, K. L., DiBlasio-Smith, E. A., McCoy, J. M., and LaVallie, E. R. (1995) *Biotechnology (NY)*, **13**, 982-987.
12. Voza, L. A., Wittwer, L., Higgins, D. R., Purcell, T. J., Bergseid, M., Collins-Racie, L. A., LaVallie, E. R., and Hoeffler, J. P. (1996) *Biotechnology (NY)*, **14**, 77-81.
13. Svetina, M., Kravec, N., Gaberc-Porekar, V., and Komel, R. (2000) *J. Biotechnol.*, **76**, 245-251.
14. Choi, S. I., Song, H. W., Moon, J. W., and Seong, B. L. (2001) *Biotechnol. Bioeng.*, **75**, 718-724.
15. Lu, D., Futterer, K., Korolev, S., Zheng, X., Tan, K., Waksman, G., and Sadler, J. E. (1999) *J. Mol. Biol.*, **292**, 361-373.
16. Gasparian, M. E., Ostapchenko, V. G., Schulga, A. A., Dolgikh, D. A., and Kirpichnikov, M. P. (2003) *Protein. Exp. Purif.*, **31**, 133-139.
17. Grant, D. A., and Hermon-Taylor, J. (1979) *Biochim. Biophys. Acta*, **567**, 207-215.
18. Green, G. D., and Shaw, E. (1979) *Analyt. Biochem.*, **93**, 223-226.
19. Schagger, H., and von Jagow, G. (1987) *Analyt. Biochem.*, **166**, 368-379.
20. Bradford, M. M. (1976) *Analyt. Biochem.*, **72**, 248-254.
21. Lu, D., Yuan, X., Zheng, X., and Sadler, J. E. (1997) *J. Biol. Chem.*, **272**, 31293-31300.
22. Rose, T., and Di Cera, E. (2002) *J. Biol. Chem.*, **277**, 19243-19246.
23. Matsushima, M., Ichinose, M., Yahagi, N., Tsukada-Kato, S., Miki, K., Omata, M., Kim, Y. T., Ito, H., Takahashi, T., Sakurai, Y., Tsuchiya, Y., Athauda, S. B., Inoue, H., and Takahashi, K. (1999) *J. Biochem. (Tokyo)*, **125**, 947-951.
24. Chen, J. M., Kukor, Z., Le Marechal, C., Toth, M., Tsakiris, L., Raguene, O., Ferec, C., and Sahin-Toth, M. (2003) *Mol. Biol. Evol.*, **20**, 1767-1777.
25. Sreedharan, S. K., Verma, C., Caves, L. S., Brocklehurst, S. M., Gharbia, S. E., Shah, H. N., and Brocklehurst, K. (1996) *Biochem. J.*, **316**, 777-786.
26. Varughese, K. I., Ahmed, F. R., Carey, P. R., Hasnain, S., Huber, C. P., and Storer, A. C. (1989) *Biochemistry*, **28**, 1330-1332.