

Structure of the K82 Capsular Polysaccharide from *Acinetobacter baumannii* LUH5534 Containing a D-Galactose 4,6-Pyruvic Acid Acetal

A. A. Kasimova^{1,2,a*}, J. J. Kenyon^{3,4,b}, N. P. Arbatsky¹, A. S. Shashkov¹,
A. V. Popova^{5,6,c}, Y. A. Knirel^{1,d}, and R. M. Hall^{3,7,e}

¹Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 119991 Moscow, Russia

²Higher Chemical College of the Russian Academy of Sciences, D. I. Mendeleev University
of Chemical Technology of Russia, 125047 Moscow, Russia

³School of Molecular Bioscience, The University of Sydney, Sydney, NSW 2006, Australia

⁴Institute of Health and Biomedical Innovation, School of Biomedical Sciences, Faculty of Health,
Queensland University of Technology, Brisbane, QLD 4059, Australia

⁵Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Moscow Region, Russia

⁶State Research Center for Applied Microbiology and Biotechnology, 142279 Obolensk, Moscow Region, Russia

⁷School of Life and Environmental Sciences, The University of Sydney, Sydney, NSW 2006, Australia

^ae-mail: nastia-kasimova979797@mail.ru

^be-mail: johanna.kenyon@qut.edu.au

^ce-mail: popova_nastya86@mail.ru

^de-mail: yknirel@gmail.com

^ee-mail: ruth.hall@sydney.edu.au

Received January 24, 2018

Revision received March 1, 2018

Abstract—Type K82 capsular polysaccharide (CPS) was isolated from *Acinetobacter baumannii* LUH5534. The structure of a linear tetrasaccharide repeating unit of the CPS was established by sugar analysis along with one- and two-dimensional ¹H and ¹³C NMR spectroscopy. Proteins encoded by the KL82 capsule gene cluster in the genome of LUH5534 were assigned to roles in the synthesis of the K82 CPS. In particular, functions were assigned to two new glycosyltransferases (Gtr152 and Gtr153) and a novel pyruvyltransferase, Ptr5, responsible for the synthesis of D-galactose 4,6-(R)-pyruvic acid acetal.

DOI: 10.1134/S0006297918070064

Keywords: *Acinetobacter baumannii*, capsular polysaccharide structure, pyruvic acid acetal, K locus, genetics of capsule biosynthesis

Acinetobacter baumannii has become one of the most widespread agents causing nosocomial infections. Currently, the majority of *A. baumannii* isolates display resistance to almost all therapeutically suitable antibiotics [1].

A thick polysaccharide capsule (CPS) surrounds the *A. baumannii* cell and protects the bacterium from the

action of immune system components, as well as disinfectants, desiccation, and some antimicrobial compounds. The CPS is composed of many oligosaccharide repeats (K units) and is characterized by high structural diversity mainly due to variability of the gene content at the chromosomal K locus (KL) driving the CPS biosynthesis. To date, more than 120 various gene clusters at the K locus have been recognized ([2, 3]; J. J. Kenyon, unpublished data). The chemical structures for about 40 different *A. baumannii* CPSs have been established ([4-8] and references cited in [4]) and form the basis for classification of strains of these bacteria. Most of these structures are consistent with putative functions of CPS synthesis genes located at the K locus.

Abbreviations: COSY, correlation spectroscopy; CPS, capsular polysaccharide; HMBC, heteronuclear multiple-bond correlation; HSQC, heteronuclear single-quantum coherence; ROESY, rotating-frame nuclear Overhauser effect spectroscopy; TOCSY, total correlation spectroscopy; Und-P, undecaprenyl phosphate.

* To whom correspondence should be addressed.




Fig. 1. ^{13}C NMR spectrum of the K82 CPS from *A. baumannii* LUH5534. Numbers refer to carbons in Pyr and sugar residues denoted by letters as shown in Table 1 and Fig. 2.

In this work, we report on the structure and gene cluster of the CPS that is specific for *A. baumannii* LUH5534. The CPS gene cluster originally designated PSgc3 has been sequenced and genes have been annotated by comparison with sequences in available databases [3]. Here, we re-annotated this gene cluster using the established nomenclature system for *A. baumannii* CPS [2] and rename it KL82. Accordingly, the CPS of *A. baumannii* LUH5534 was assigned to the K82 type.

MATERIALS AND METHODS

Cultivation of bacteria. *Acinetobacter baumannii* strain LUH5534 was cultivated in 2×TY media for 24 h. Cells were harvested by centrifugation (10,000g, 20 min), washed with phosphate-buffered saline (pH 7.4), suspended in a 7 : 3 acetone–water mixture (v/v), precipitated, and dried.

Isolation of capsular polysaccharide. CPS was isolated by phenol–water extraction [9] of bacterial cells (1.12 g).

Table 1. ^1H and ^{13}C NMR chemical shifts (δ , ppm) in K82 CPS from *A. baumannii* LUH5534

Residue	C1 <i>H1</i>	C2 <i>H2</i>	C3 <i>H3</i>	C4 <i>H4</i>	C5 <i>H5</i>	C6 <i>H6 (6a,6b)</i>
CPS						
→3)-β-D-GalpNAc-(1→ A	103.4 <i>4.76</i>	52.6 <i>4.03</i>	82.0 <i>3.82</i>	69.5 <i>4.14</i>	76.1 <i>3.65</i>	61.7 <i>3.82</i>
→3)-β-D-Galp-(1→ B	106.0 <i>4.47</i>	70.5 <i>3.63</i>	78.7 <i>3.66</i>	66.4 <i>4.04</i>	76.1 <i>3.59</i>	62.4 <i>3.71; 3.76</i>
→3)-α-D-GlcpNAc-(1→ C	95.7 <i>5.01</i>	53.0 <i>4.13</i>	80.3 <i>4.06</i>	68.8 <i>3.70</i>	72.8 <i>4.01</i>	62.5 <i>3.75; 3.83</i>
→2,4,6)-β-D-Galp-(1→ D	100.5 <i>4.64</i>	79.2 <i>3.75</i>	73.6 <i>3.75</i>	72.7 <i>4.13</i>	67.1 <i>3.55</i>	66.2 <i>3.90; 4.02</i>
Pyr	175.2	101.2	26.3 <i>1.50</i>			
MPS						
→3)-β-D-GalpNAc-(1→ A	103.1 <i>4.77</i>	52.6 <i>4.03</i>	82.2 <i>3.82</i>	69.6 <i>4.15</i>	76.1 <i>3.64</i>	61.6 <i>3.82</i>
→3)-β-D-Galp-(1→ B	106.1 <i>4.47</i>	70.4 <i>3.62</i>	78.6 <i>3.67</i>	66.3 <i>4.06</i>	76.1 <i>3.60</i>	62.3 <i>3.75</i>
→3)-α-D-GlcpNAc-(1→ C	95.5 <i>5.02</i>	53.6 <i>4.11</i>	80.1 <i>4.08</i>	69.2 <i>3.66</i>	69.9 <i>4.00</i>	62.4 <i>3.73; 3.82</i>
→2)-β-D-Galp-(1→ D	101.8 <i>4.59</i>	79.5 <i>3.66</i>	75.0 <i>3.69</i>	70.1 <i>3.86</i>	76.4 <i>3.69</i>	62.4 <i>3.73</i>

Notes: ^1H NMR chemical shifts are italicized. Chemical shifts for N-acetyl groups are: δ_{C} 23.4–23.6 (Me) and 175.2–176.0 (CO), δ_{H} 2.01–2.13. MPS, modified polysaccharide

Table 2. Correlations for H1 and C1 in the two-dimensional ^1H , ^1H ROESY and ^1H , ^{13}C HMBC spectra of the K82 CPS from *A. baumannii* LUH55534

Atom in sugar residue (δ)	Correlations to atoms in sugar residues (δ)	
	^1H , ^1H ROESY	^1H , ^{13}C HMBC
A H1 (4.76)	D H2 (3.75), A H3 (3.82), A H5 (3.65)	D C-2 (79.2)
A C1 (103.4)		D H2 (3.75), A H2 (4.03), A H5 (3.65)
B H1 (4.47)	A H3 (3.82), B H2 (3.63), B H3 (3.66), B H5 (3.59)	A C3 (82.0)
B C1 (106.0)		A H3 (3.82), B H2 (3.63), B H5 (3.59)
C H1 (5.01)	B H2 (3.63), B H3 (3.66), B H4 (4.04)	B C3 (78.7), C C3 (80.3), C C5 (72.8)
C C1 (95.7)		B H3 (3.66)
D H1 (4.64)	C H2 (4.13; w), C H3 (4.06), C H4 (3.70; w), D H3 (3.75), D H5 (3.55)	C C3 (80.3)
D C1 (100.5)		C H3 (4.06), D H2 (3.75), D H5 (3.55)

Note: w, weak.

(*R*)-configured D-galactose 4,6-pyruvic acid acetal. Earlier, a similar acetal but on a D-GalpNAc residue was reported in the K4 CPS from *A. baumannii* D78 [12].

The KL82 gene cluster in *A. baumannii* LUH5534 (GenBank accession number KC526908) between con-

served genes *gna* and *galU* contains genes involved specifically with the synthesis of the K82 CPS (Fig. 3).

The *itrA2* gene in the KL82 locus encodes an initiating transferase, ItrA2 (WeeH in GenPept, accession number AHB32565.1), which is 99% identical to ItrA2 of *A. baumannii* KL2 (AGK44809.1 in GenPept) shown to use D-GalNAc as the initiating sugar in the K-unit assembly [13, 14]. ItrA2 in KL82 is 99% identical to ItrA2 of *A. baumannii* ATCC 17978 (KL3). Hence, D-GalNAc is the first sugar of the K82 unit, and the Wzy polymerase (AHB32563.1 in GenPept) would catalyze formation of the β -D-GalpNAc-(1 \rightarrow 3)-D-Galp linkage between the K units (Fig. 2).

Glycosyltransferase Gtr5 encoded by KL82 (WafH in GenPept, accession number AHB32564.1) is 95% identical to Gtr5a from *A. baumannii* KL2 (GenPept accession number AHM95430.1), which is responsible for formation of the β -D-Galp-(1 \rightarrow 3)-D-GalpNAc linkage in the K2 CPS [15]. This linkage is the first in the K82 unit, and Gtr5 was assigned accordingly.

Two further glycosyltransferase genes, *gtr152* and *gtr153*, were also identified in KL82 (Fig. 3) and are evidently responsible for adding the next two sugar residues to complete the assembly of the tetrasaccharide K unit (Fig. 2). Gtr153 (WafK in GenPept, accession number AHB32562.1) is 54% identical to Gtr58 from *A. baumannii* KL27 (ALL34866.1 in GenPept) that was previously predicted to form an α -D-GlcpNAc-(1 \rightarrow 3)-D-Galp linkage [16]. Accordingly, Gtr153 would catalyze this linkage, which also is present in the K82 CPS, and Gtr152 (WafJ in GenPept, accession number AHB32561.1) was assigned to the last β -D-Galp-(1 \rightarrow 3)- α -D-GlcpNAc linkage (Fig. 2).

The product of the gene located immediately downstream of the *wzx* gene was identified as a pyruvyl transferase belonging to protein family (Pfam) PF04230 and was named Ptr5 (WafI in GenPept, accession number AHB32560.1). Ptr5 is not significantly related to Ptr1 from *A. baumannii* KL4 (GenPept accession number JN409449.3) putatively involved with formation of a D-GalNAc4,6Pyr [16], though it would be responsible for




Fig. 3. Comparison of the *A. baumannii* KL82 and KL124 capsule biosynthesis gene clusters. Shading between gene clusters indicates shared regions of >95% nucleotide sequence identity.

the similar attachment of pyruvic acid to the Gal residue (D) in the K82 CPS.

The generally conserved *galU-ugd-gpi-gne1-pgm* arrangement in *A. baumannii* is interrupted in KL82 by a partial copy of ISAbal3 (138-1016 of 1039 bp) and a gene encoding a putative acyltransferase (GenPept accession number AHB32571.1 previously annotated as CgmA) that belongs to Pfam PF01757. The first 142 amino acids of the acyltransferase (201 a.a. totally) is 82% identical to Atr5 (142 a.a.) from *A. baumannii* KL4 (GenPept accession number ACJ39541.2), and neither appears to modify the K4 [12] or K82 CPS. An additional gene cluster KL124 differs from KL82 only in this region (Fig. 3).

Finally, the occurrence in KL82 of genes for flippase Wzx and polymerase Wzy indicated that, as for all other known CPSs of *A. baumannii*, the K82 CPS is synthesized by the Wzx/Wzy-dependent pathway. Therefore, the content of the KL82 gene cluster and predicted gene functions are consistent with the K82 CPS structure established in this work.

Acknowledgments

This work was supported by the Russian Foundation for Basic Research (project No. 17-04-01254) and the National Health and Medical Research Council (project No. 1026189).

REFERENCES

- World Health Organization (2017) Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics (<http://www.who.int/medicines/publications/global-priority-list-antibiotic-resistant-bacteria/en/>).
- Kenyon, J. J., and Hall, R. M. (2013) Variation in the complex carbohydrate biosynthesis loci of *Acinetobacter baumannii* genomes, *PLoS One*, **8**, e62160.
- Hu, D., Liu, B., Dijkshoorn, L., Wang, L., and Reeves, P. R. (2013) Diversity in the major polysaccharide antigen of *Acinetobacter baumannii* assessed by DNA sequencing, and development of a molecular serotyping scheme, *PLoS One*, **8**, e70329.
- Kenyon, J. J., Kasimova, A. A., Shneider, M. M., Shashkov, A. S., Arbatsky, N. P., Popova, A. V., Miroshnikov, K. A., Hall, R. M., and Knirel, Y. A. (2017) The KL24 gene cluster and a genomic island encoding a Wzy polymerase contribute genes needed for synthesis of the K24 capsular polysaccharide by the multiply antibiotic resistant *Acinetobacter baumannii* isolate RCH51, *Microbiology*, **163**, 355-363.
- Kasimova, A. A., Shneider, M. M., Arbatsky, N. P., Popova, A. V., Shashkov, A. S., Miroshnikov, K. A., Balaji, V., Biswas, I., and Knirel, Y. A. (2017) Structure and gene cluster of the capsular polysaccharide of *Acinetobacter baumannii* B11911 containing 5-*N*-acetyl-7-*N*-[(*R*)-3-hydroxybutanoyl]pseud-aminic acid, *Biochemistry (Moscow)*, **82**, 483-489.
- Kenyon, J. J., Shashkov, A. S., Senchenkova, S. N., Shneider, M. M., Liu, B., Popova, A. V., Arbatsky, N. P., Miroshnikov, K. A., Wang, L., Knirel, Y. A., and Hall, R. M. (2017) *Acinetobacter baumannii* K11 and K83 capsular polysaccharides have the same 6-deoxy-L-talose-containing pentasaccharide K units but different linkages between the K units, *Int. J. Biol. Macromol.*, **103**, 648-655.
- Shashkov, A. S., Liu, B., Kenyon, J. J., Popova, A. V., Shneider, M. M., Senchenkova, S. N., Arbatsky, N. P., Miroshnikov, K. A., Wang, L., and Knirel, Y. A. (2017) Structures of the K35 and K15 capsular polysaccharides of *Acinetobacter baumannii* LUH5535 and LUH5554 containing amino and diamino uronic acids, *Carbohydr. Res.*, **448**, 28-34.
- Kenyon, J. J., Kasimova, A. A., Notaro, A., Arbatsky, N. P., Speciale, I., Shashkov, A. S., De Castro, C., Hall, R. M., and Knirel, Y. A. (2017) *Acinetobacter baumannii* K13 and K73 capsular polysaccharides differ only in K-unit side branches of novel non-2-ulosonic acids: di-*N*-acetylated forms of either acinetaminic acid or 8-epiacinetaminic acid, *Carbohydr. Res.*, **452**, 149-155.
- Westphal, O., and Jann, K. (1965) Bacterial lipopolysaccharides. Extraction with phenol-water and further applications of the procedure, *Methods Carbohydr. Chem.*, **5**, 83-91.
- Lipkind, G. M., Shashkov, A. S., Knirel, Y. A., Vinogradov, E. V., and Kochetkov, N. K. (1988) A computer-assisted structural analysis of regular polysaccharides on the basis of ¹³C NMR data, *Carbohydr. Res.*, **175**, 59-75.
- Garegg, P. J., Lindberg, B., and Kvarnstrom, I. (1980) Preparation and NMR studies of pyruvic acid and related acetals of pyranosides: configuration at the acetal carbon atoms, *Carbohydr. Res.*, **77**, 71-78.
- Kenyon, J. J., Speciale, I., Hall, R. M., and De Castro, C. (2016) Structure of repeating unit of the capsular polysaccharide from *Acinetobacter baumannii* D78 and assignment of the K4 gene cluster, *Carbohydr. Res.*, **408**, 12-17.
- Iwashkiw, J. A., Seper, A., Weber, B. S., Scott, N. E., Vinogradov, E. V., Stratilo, C., Reiz, B., Cordwell, S. J., Whittall, R., Schild, S., and Feldman, M. F. (2012) Identification of a general O-linked protein glycosylation system in *Acinetobacter baumannii* and its role in virulence and biofilm formation, *PLoS Pathog.*, **8**, e1002758.
- Lees-Miller, R. G., Iwashkiw, J. A., Scott, N. E., Seper, A., Vinogradov, E., Schild, S., and Feldman, M. F. (2013) A common pathway for O-linked protein glycosylation and synthesis of capsule in *Acinetobacter baumannii*, *Mol. Microbiol.*, **89**, 816-830.
- Kenyon, J. J., Marzaioli, A. M., Hall, R. M., and De Castro, C. (2014) Structure of the K2 capsule associated with the KL2 gene cluster of *Acinetobacter baumannii*, *Glycobiology*, **24**, 554-563.
- Shashkov, A. S., Kenyon, J. J., Senchenkova, S. N., Shneider, M. M., Popova, A. V., Arbatsky, N. P., Miroshnikov, K. A., Volozhantsev, N. V., Hall, R. M., and Knirel, Y. A. (2016) *Acinetobacter baumannii* K27 and K44 capsular polysaccharides have the same K unit but different structures due to the presence of distinct *wzy* genes in otherwise closely related K gene clusters, *Glycobiology*, **26**, 501-508.